



TITLE:

A unified family of $P_{\{\mathrm{J}\}}$ -hierarchies ($J=I, II, IV, 34$) with a large parameter (Algebraic analytic methods in complex partial differential equations)

AUTHOR(S):

Umeta, Yoko

CITATION:

Umeta, Yoko. A unified family of $P_{\{\mathrm{J}\}}$ -hierarchies ($J=I, II, IV, 34$) with a large parameter (Algebraic analytic methods in complex partial differential equations). 数理解析研究所講究録 2017, 2020: 92-95

ISSUE DATE:

2017-04

URL:

<http://hdl.handle.net/2433/231742>

RIGHT:

A unified family of P_J -hierarchies ($J=I, II, IV, 34$) with a large parameter

By

Yoko UMETA*

Abstract

The purpose of this note is to give a unified family of P_J -hierarchies ($J=I, II, IV, 34$) with a large parameter. This note is a short summary of papers [18] and [19].

§ 1. A unified family of P_J -hierarchies ($J=I, II, IV, 34$)

The P_I , P_{II} , P_{IV} and P_{34} -hierarchies were studied by Kudryashov ([11],[12]), Gordoa and Pickering ([4]), Shimomura ([13],[14]), Gordoa, Joshi and Pickering ([5]), Clarkson, Joshi and Pickering ([3]). To establish connection formulas for solutions of the higher order Painlevé equations is one of important subjects in algebraic analysis of singular perturbation theory. In the series of papers by Kawai, Koike, Nishikawa and Takei, they introduced a large parameter η to P_J -hierarchies ($J = I, II, IV, 34$) ([6], [9], [10]) and many important results have been established from a view point of the exact WKB analysis. (See [6], [7], [8], [15], [16], [17] and etc). In what follows, we give a unified family of P_J -hierarchies ($J = I, II, IV, 34$) with a large parameter η .

Let m be an arbitrary natural number. Let U , V and C denote generating functions of unknown functions u_k , v_k ($k = 1, 2, \dots, m$) and constants c_k as follows.

$$U(\theta) := \sum_{k=1}^{m+1} u_k \theta^k, \quad V(\theta) := \sum_{k=1}^{m+1} v_k \theta^k, \quad C(\theta) := \sum_{k=1}^m c_k \theta^k.$$

Here θ denotes an independent variable, u_{m+1} and v_{m+1} are arbitrary holomorphic functions of t . Throughout the note, the notation $A \equiv B$ means that $A - B$ is zero

2010 Mathematics Subject Classification(s): Primary 34E20; Secondary 34M40.

Key Words: exact WKB analysis, Painlevé hierarchies, instanton-type solutions

This work was supported by JSPS KAKENHI Grant No.15K17557.

*Department of Mathematical Sciences, Yamaguchi University, Yamaguchi City, Yamaguchi Pref. 753-8512, Japan.

modulo θ^{m+2} . Let us consider the system of non-linear ordinary differential equations with a large parameter η for these generating functions:

$$(1.1) \quad \eta^{-1} \frac{d}{dt} \begin{pmatrix} U\theta \\ V\theta \end{pmatrix} \equiv \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} \times (1-U) + \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \frac{\partial H}{\partial U} \\ \frac{\partial H}{\partial V} \end{pmatrix} + \begin{pmatrix} 0 \\ \frac{H(U,V)}{1-U} \end{pmatrix},$$

where $H(U, V)$ is a polynomial in U and V with arbitrary complex constants p_i of the following form

$$H(U, V) := (p_1 U^2 + p_2 V^2)\theta + p_3 UV + p_4 CU + p_5 CV + p_6 U + p_7 V + p_8 C + p_9,$$

and f_1, f_2 are defined by

$$\begin{aligned} f_1 &:= p_7 + (\alpha u_1 + p_5 c_1)\theta + (y_1 + (y_1 u_1 + y_2)\theta)\theta^m, \\ f_2 &:= -\beta - (2\beta u_1 + \alpha v_1 + \varepsilon c_1)\theta + (z_1 + (2z_1 u_1 - y_1 v_1 + z_2)\theta)\theta^m. \end{aligned}$$

Here y_i, z_i are arbitrary holomorphic functions of t and $\alpha, \beta, \varepsilon$ are given by

$$\alpha := p_3 + p_7, \quad \beta := p_6 + p_9 \quad \text{and} \quad \varepsilon := p_4 + p_8,$$

respectively.

If p_i, y_i, z_i are determined as follows, then (1.1) is same as the general member $(P_J)_m$ of P_J -hierarchy with η (See [19]).

- If $p_2 = -1, p_8 = 2, p_9 = 1, z_2 = 2t$, the others $= 0 \implies (P_I)_m$.
- If $p_2 = -1, p_8 = 2, p_9 = 1, z_1 = 2\gamma t (\gamma \neq 0), z_2 = 4\gamma t c_0$, the others $= 0 \implies (P_{34})_m$.
- If $p_2 = 1, p_3 = 2, p_5 = 2$, the others $= 0 \implies (P_{II})_m$.
- If $p_2 = 1, p_3 = 2, p_5 = 2, y_1 = -2\gamma t (\gamma \neq 0)$, the others $= 0 \implies (P_{IV})_m$.

§ 2. The existence of general formal solutions of (1.1)

We can apply the method given in [2] to the cases I, II:

Case I: $\alpha = p_3 + p_7 \neq 0, \quad p_2 \neq 0$.

Case II: $\alpha = p_3 + p_7 = 0, \quad \beta = p_6 + p_9 \neq 0, \quad p_2 \neq 0$

and we have the following theorem. (For more precise statements, see [19].)

Theorem 2.1. *In the cases I, II, we have formal solutions with $2m$ free parameters called instanton-type solutions for (1.1).*

§ 3. Lax pair for (1.1)

Theorem 3.1. *Let us determine p_1 , u_{m+1} and v_{m+1} of (1.1) so that they satisfy the following conditions.*

$$\begin{cases} p_1 = 0, & p_2 \neq 0, \\ \gamma \alpha \theta^{k-2} = -\alpha u'_{m+1} \theta^{m+1} + (y_1' \theta^m + y_2' \theta^{m+1}), \\ z_1' \theta^m + (z_1' u_1 - y_1' v_1 + z_2') \theta^{m+1} + (2\beta u'_{m+1} + \alpha v'_{m+1}) \theta^{m+1} + \gamma \beta \theta^{k-2} = 0. \end{cases}$$

Here u'_{m+1} denotes the derivative of u_{m+1} with respect to t , γ is a non-zero constant, and k is determined by the above conditions. Then our system (1.1) is equivalent to the compatibility condition of the following equations:

$$(I) \quad \left(\gamma \theta^k \frac{\partial}{\partial \theta} - \eta A \right) \psi(\theta, t) = 0, \quad (II) \quad \left(\frac{\partial}{\partial t} - \eta B \right) \psi(\theta, t) = 0,$$

where

$$A := \begin{pmatrix} \Delta_1 (1 - U) \theta \\ \Delta_2 & -\Delta_1 \end{pmatrix}, \quad B := \begin{pmatrix} \square_1 & 1 \\ \square_2 & -\square_1 \end{pmatrix}$$

with

$$\begin{aligned} \Delta_1 &:= -\frac{1}{2} \frac{\partial H}{\partial V} - \frac{p_3}{2} (1 - U) + \frac{1}{2} (y_1 \theta^m + y_2 \theta^{m+1}) - \frac{\alpha}{2} u_{m+1} \theta^{m+1}, \\ \Delta_2 &:= p_2 \times \left(-\frac{\partial H}{\partial U} - \frac{H(U, V)}{1 - U} - (z_1 \theta^m + (z_1 u_1 - y_1 v_1 + z_2) \theta^{m+1}) \right. \\ &\quad \left. - (2\beta u_{m+1} + \alpha v_{m+1}) \theta^{m+1} \right), \\ \square_1 &:= -\frac{1}{2\theta} (\alpha + (\alpha u_1 + p_5 c_1) \theta), \\ \square_2 &:= -\frac{p_2}{\theta} (\beta + (2\beta u_1 + \alpha v_1 + \varepsilon c_1) \theta). \end{aligned}$$

The Lax pair associated with (1.1) plays an important role in analyzing the Stoke geometry of (1.1) (See [18]).

Acknowledgments The author would like to express her sincere gratitude to Professors Yasunori Okada, Naofumi Honda and Hiroshi Yamazawa for giving the opportunity to her to take part in the conference.

References

- [1] Aoki, T., Multiple-scale analysis for higher-order Painlevé equations, *RIMS Kôkyûroku Bessatsu* **B5** (2008), 89–98.

- [2] Aoki, T., Honda, N. and Umeta, Y., On a construction of general formal solutions for equations of the first Painlevé hierarchy I, *Adv. in Math.*, **235** (2013), 496–524.
- [3] Clarkson, P. A., Joshi, N., Pickering, A., Bäcklund transformations for the second Painlevé hierarchy: a modified truncation approach, *Inverse Problems* **15** (1999) 175–187.
- [4] Gordoa, P. R. and Pickering, A., Nonisospectral scattering problems: A key to integrable hierarchies, *J. Math. Phys.*, **40** (1999), 5749–5786.
- [5] Gordoa, P. R. and Pickering, A., On a generalized 2+1 dispersive water wave hierarchy, *Publ. RIMS, Kyoto Univ.*, **37** (2001), 327–347.
- [6] Kawai, T., Koike, T., Nishikawa, Y. and Takei, Y., On the Stokes geometry of higher order Painlevé equations, *Astérisque* **297** (2004), 117–166.
- [7] Kawai, T. and Takei, Y., WKB analysis of higher order Painlevé equations with a large parameter - Local reduction of 0-parameter solutions for Painlevé hierarchies (P_J) ($J = \text{I, II-1 or II-2}$), *Adv. Math.*, **203** (2006), 636–672.
- [8] ———, WKB analysis of higher order Painlevé equations with a large parameter. II. Structure theorem for instanton-type solutions of $(P_J)_m$ ($J = \text{I, 34, II-2 or IV}$) near a simple P -turning point of the first kind, *Pub. RIMS, Kyoto Univ.* **47** (2011), 153–219.
- [9] Koike, T., On the Hamiltonian structures of the second and the forth Painlevé hierarchies and degenerate Garnier systems, *RIMS Kôkyûroku Bessatsu* **B2** (2007), 99–127.
- [10] ———, On new expressions of the Painlevé hierarchies, *RIMS Kôkyûroku Bessatsu* **B5** (2008), 153–198.
- [11] Kudryashov, N. A., The first and second Painlevé equations of higher order and some relations between them, *Phys. Lett. A*, **224** (1997), 353–360.
- [12] Kudryashov, N. A. and Soukharev, M. B., Uniformization and transcendence of solutions for the first and second Painlevé hierarchies, *Phys. Lett. A*, **237** (1998), 206–216.
- [13] S. Shimomura, Painlevé properties of a degenerate Garnier system of $(9/2)$ -type and of a certain fourth order non-linear ordinary differential equation, *Ann. Scuola Norm. Sup. Pisa*, **29** (2000), 1–17.
- [14] ———, A certain expression of the first Painlevé hierarchy, *Proc. Japan Acad., Ser. A*, **80** (2004), 105–109.
- [15] Takei, Y., An explicit description of the connection formula for the first Painlevé equation, Toward the Exact WKB Analysis of Differential Equations, Linear or Non-Linear, *Kyoto University Press* (2000), 271–296.
- [16] ———, Toward the exact WKB analysis for instanton-type solutions of Painlevé hierarchies, *RIMS Kôkyûroku Bessatsu* **B2** (2007), 247–260.
- [17] ———, Instanton-type formal solutions for the first Painlevé hierarchy, in: *Algebraic Analysis of Differential Equations*, Springer-Verlag (2008), 307–319.
- [18] Umeta, Y., On the Stokes geometry of a unified family of P_J -hierarchies ($J = \text{I, II, IV, 34}$), in preparation.
- [19] ———, General formal solutions for a unified family of P_J -hierarchies ($J = \text{I, II, IV, 34}$), in preparation.